



Organic Nitrogen (N)	1%
Organic Carbon (C)	12%
Potassium Oxide (K2O)	6%
Betaine	1%

ALGAFLUID®

VEGETAL BIOSTIMULANT FOR ORGANIC FARMING

ALGAFLUID, entirely obtained from vegetal constituents has a biostimulant action improving the plants growth especially in case of difficult climatic conditions. ALGAFLUID helps nutrients intake and it's promptly absorbed. It moves rapidly from the leaves surface to the cells increasing the metabolic activity of the plant.

DOSAGE AND USAGE PROCEDURES

FOLIAR APPLICATIONS	150-200 grams per 100 lt water
FERTIGATION	5-10 Kg/Ha







Tanks Kg. 6-30



AMINOSLOW®

LIQUID ORGANO-MINERAL FERTILIZER WITH SLOW-RELEASE NITROGEN

Organo-mineral fertilizer with a high content of natural organic and synthesised (methylenurea) nitrogen. It allows useful nitrogen nutrition in all vegetative phases. The particular ratio between the various forms of nitrogen provides continuous nutrition for all crops. The presence of methylenurea (slow release nitrogen) and amino acids (the organic part of the product) gives this formulation unique nutritive characteristics whereby the Nitrogen is entirely and gradually available. This is especially beneficial during periods of maximum growth and in conditions of climatic stress.

DOSAGE AND USAGE PROCEDURES

Citrus - Top Fruit	150 g/Hl (foliar) 20-40 Kg/Ha (fertigation)
Grapes	100 g/Hl (foliar) 30-50 Kg/Ha (fertigation)
Field and Greenhouse vegetables	100 g/Hl (foliar) 30 Kg/Ha (fertigation)
Floriculture	100 g/Hl (foliar) 30 Kg/Ha (fertigation)



COMPOSITION

Total Nitrogen (N)12%
Ureic Nitrogen (N)6%
Organic Nitrogen (N)6%
Methylene-Urea Nitrogen (N)4%
Biological Organic Carbon (C)25%



Kg. 6-30





Total Nitrogen (N)9%	, o
Organic Nitrogen (N)9%	
Biological Organic Carbon (C)25%	ó

AMINOGRAM (gr/100 gr di amino acid)

Aspartic Acid	5 70
Glutamic Acid	
Alanine	
Arginine	
Cysteine	
Phenylalanine	
Glycine	25,31
Hydroxyproline	8,18
Isoleucine	1,48
Istidine	
Leucine	
Lysine	
Methionine	0.74
Proline	
Serine	
Tyrosine	,
Threonine	
Tryptophan	
Valine	
vaiiiie	2,40





Kg. 1



Tanks Ka. 6-30

AMINOSPRAY

ORGANIC LIQUID FERTILIZER WITH AMINO-ACIDS FROM ENZYMATIC HYDROLYSIS

A latest generation organic fertilizer made of low molecular weight amino acids, peptides and peptones formulated for repeated foliar treatments to all crops, horticultural and arboreal. The amino acids contained are indispensable for the crop's life, not only as nutrients but also as catalysts of enzymatic activities essential for plant metabolism. Proteins are a vital constituent of animal and plant life: in the latter being indispensable for vegetative growth, flowering and fruit forma-tion. Starting with Nitrogen, water and air pho-tochemical processes convert these raw materials initially into amino acids: subsequently peptides and proteins are formed. This requires time and significant use of the crop's energy intake. AMINOSPRAY, supplying readily absor-bed peptides and peptones accelerates protein formation and increases the productive capacity of the treated crops. Nutrient penetration into the tissues is also enhanced reducing stress due to climatic factors and/or pesticide treatments.

The product can be used both as a foliar application and in fertigation systems.

DOSAGE AND USAGE PROCEDURES

Top Fruit - Grapes - Olive	300 g/HI
Floriculture and Greenhouse vegetables	300 g/Hl
Field vegetables	250 g/Hl
Cereal	4-5 Kg/Ha



BIOCAL

CALCIUM BASED LIQUID FERTILIZER

BIOCAL is formulated to ensure a high nutrition in calcium coformulated with proteininc extracts helping the crops to develop their natural defence system. The plants are more resistant and promptly overcome all the types of stress. The high content in calcium improves the colour and texture of the fruits increasing production yields.

DOSAGE AND USAGE PROCEDURES

Fruit trees Vegetables	150-200 g/100 lt water (foliar)
---------------------------	---------------------------------



COMPOSITION


Calcium Oxide (CaO).....12% water soluble coformulated with proteininc extracts







Tanks Kg. 6







Boron (B)11% water soluble (as ethanolamine salt)

BOROFLUID

LIQUID FERTILIZER CONTAINING BORON AS ETHANOLAMINE SALT

BOROFLUID is formulated to prevent and cure Boron shortages caused by imbalances and deficiencies in the assimilation of this element. Boron is essential for the crop's development and deficiency leads to decreased production, poor vegetative growth and loss of quality in the fruits. Boron moves poorly inside the crop and therefore has to be supplied as a preventative treatment.

BOROFLUID can be used both as a foliar application and in fertigation systems.

DOSAGE AND USAGE PROCEDURES

Citrus - Top Fruit	100 g/Hl (foliar)
Grapes	10 Kg/Ha (fertigation)
Olive	100 g/Hl (foliar) 10 Kg/Ha (fertigation)
Field	50-100 g/Hl (foliar)
vegetables	5-10 Kg/Ha (fertigation)
Floriculture	50-100 g/Hl (foliar) 5-10 Kg/Ha (fertigation)





Bottles Kg. 1 Tanks Kg. 6



EMOFLUID®

LIQUID ORGANO-MINERAL FERTILIZER WITH NITROGEN FROM BLOOD MEAL AND METHYLENE UREA

EMOFLUID is a very active product intensifying the activity of the bacterial flora in the soil. This gives immediate effects on young roots, stimulating growth and nutrient intake potential. For these reasons it is recommended for the stimulation of seed germination and root development. The product is also particularly effective on vegetables and arboreal crops under climatic stress conditions due to various causes (sudden temperature change, damage caused by hail or other meteorological phenomena).

DOSAGE AND USAGE PROCEDURES (fertigation)

Citrus - Top Fruit	40-100 Kg/Ha
Grapes	80-160 Kg/Ha
Field and Greenhouse vegetable	50-100 Kg/Ha
Floriculture	50-100 Kg/Ha



COMPOSITION

Total Nitrogen (N)17%
Organic Nitrogen (N)2%
Methylene Urea Nitrogen (N)15%
Biological Organic Carbon (C)8%









Chelated Iron (Fe)6% water soluble
(MAXIRON) Ortho-Ortho EDDHA min4,8%
(EXTRAIRON) Ortho-Ortho EDDHA min. 4,2%
(FERROFORTE) Ortho-Ortho EDDHA min3,6%
Chelating agent: EDDHA
Chelate stability pH range3-11





Bags Carton case Kg. 1-2,5-5 Kg. 1

FERRO FORTE MAXIRON EXTRAIRON

EDDHA CHELATE IRON MICRO-GRANULES

Soluble microgranular formulation of Iron chelated with EDDHA, ethylendiamino-di (ohydroxyphenylacetic) acid. The main characteristic of this product is the presence of the stable ortho-ortho form of the EDDHA chelating agent. Iron chelated with this isomer, once distributed in the soil, is released to the plant even in the case of soils with high pH values becoming essential for the treatment of iron chlorosis, especially in calcareous soils where iron salts are precipitated and locked in the soil, unavailable to the plant.

Products applications are suggested all the times that the plants show symptoms of yellowing between the "veins" of the leaves, especially noticeable in young shoots.

Because of this capacity to improve photosynthesis, the chelated iron promotes blooming and better fruit development.

DOSAGE AND USAGE PROCEDURES

Citrus - Top Fruit Olive	20-50 g/plant (preventive) 50-100 g/plant (curative)
Grapes	20-30 g/plant (preventive) 30-75 g/plant (curative)
Strawberry - Turf Vegetable crops	1-2 Kg/1000 mt







GLUCOMPOST MIX

LIQUID FOLIAR FERTILIZER WITH COMPLEXED MICRONUTRIENTS

GLUCOMPOST MIX is a liquid fertilizer designed to reduce nutrient deficiencies by inducing rapid absorption of the micronutrient content. Complexes with high molecular weight organic acid are, in fact, easily absorbed by the crop quickly transferring nutrients into the body of the plant.

DOSAGE AND USAGE PROCEDURES

Citrus - Top Fruit	300 g/HI (foliar)
Grapes	5-10 Kg/Ha (fertigation)
Greenhouse	200 g/HI (foliar)
vegetables	5-10 Kg/Ha (fertigation)
Field	300 g/Hl (foliar)
vegetables	5-10 Kg/Ha (fertigation)
Floriculture	200 g/HI (foliar) 5-10 Kg/Ha (fertigation)



COMPOSITION

Boron (B) water soluble1% Iron (Fe) EDTA chelated1% water soluble
Manganese (Mn) EDTA chelated2% water soluble
Molybdenum (Mo)0,15% water soluble
Zinc (Zn) EDTA chelated1% water soluble





Bottles Kg. 1

Tanks Kg. 6





Total Organic Nitrogen (N)	.8%
Biological Organic Carbon (C)2	7%

DOSAGE AND USAGE PROCEDURES (fertigation)

Citrus	40-100 Kg/Ha
Top Fruit - Grapes	50-100 Kg/Ha
Field and Greenhouse vegetables	30-60 Kg/Ha
Floriculture	20-40 Kg/Ha



Ka. 1200







Tanks Ka. 30



LIQUID FERTILIZER CONTAINING ORGANIC NITROGEN FROM EPITHELIUM

HERGOTON consists of low to medium molecular weight amino acids derived from the hydrolysis of animal proteins. These amino acids are rapidly and completely absorbed by the root system. HERGOTON cannot, therefore, be considered a traditional nitrogen fertilizer but as a natural and stable mixture of complex structures that represent the essential precursors for the construction of peptides, proteins and enzymes. Applying this product accelerates crop metabolism, supplying directly, in a readily available form, all those constituents that would otherwise need to be synthesised "afresh" from inorganic nitrogen. The effects obtained with root application are as follows:

- ·Increase in the activity of the soil microflora.
- ·Improvement in the availability of metallic ions thanks to the chelating action of amino acids.
- · Anti-stress effect following meteorological or parasitic damage.
- ·Inducement to biological and biochemical functions which determine vegetative growth and increased harvest.
- · Possibility of supplying organic nitrogen via fertigation.
- ·Increased root system development and activity.



LEAFEED®

ORGANIC FERTILIZER WITH LOW MOLECULAR WEIGHT AMINO-ACIDS AND NATURAL AUXINS

LEAFEED is a natural organic fertilizer containing low molecular weight amino acids, vitamins and auxins easily absorbed by all crops both arboreal and herbaceous. Stimulates growth using the effect of natural enzymes. Particularly suitable in cases of stress due to excess heat or frost. Increases the absorption and efficiency of co-applied fertilizers. The composition of **LEAFEED** promotes stomatal opening thus improving foliar intake. **LEAFEED** can be used both as a foliar application and in fertigation systems.

DOSAGE AND USAGE PROCEDURES

Citrus - Top Fruit	250 g/Hl (foliar)
Grapes	10 Kg/Ha (fertigation)
Greenhouse vegetables	200 g/HI (foliar)
Floriculture	10 Kg/Ha (fertigation)
Field vegetables	250 g/HI (foliar) 10 Kg/Ha (fertigation)

IN CASE OF FOLIAR APPLICATION, MIXTURES WITH COPPER BASED PRODUCTS TO BE AVOIDED





Bottles Kg. 1 Tanks Kg. 6



COMPOSITION

Total Organic Nitrogen (N)6%
Organic Nitrogen (N)6% water soluble
Biological Organic Carbon (C)18%

AMINOGRAM (gr/100 gr di amino acid)

Aspartic Acid	6.20
Glutamic Acid	
Alanine	
Arginine	
Cysteine	
Phenylalanine	
Glycine	25,90
Hydroxyproline	
Isoleucine	
Istidine	
Leucine	
Lysine	5,10
Methionine	1,05
Proline	
Serine	2,00
Tyrosine	1,90
Threonine	
Tryptophan	0,85
Valine	2,98

Dosage always depends on different factors (plant's age, growth stage, climatic conditions, temperature, etc). The product displays good miscibility and compatibility, however, we advise preliminary tests and treatment of a few plants before extending to the whole area. MIXTURES WITH COPPER BASED PRODUCTS TO BE AVOIDED.





Potassium Oxide (K2O)	19%
Organic Nitrogen (N)	1%
Betaine	0,1%
Mannitol	4%
Organic Carbon (C)	20%



Kg. 1



ORGANIC FERTILIZER WITH
WATER-SOLUBLE EXTRACT FROM BROWN
ALGAE (ASCOPHILLUM NODOSUM)

SEAGROWING is a natural product composed of organic substances extracted through a physico-mechanical process from brown algae (Ascophyllum Nodosum). This method preserves the activity of natural constituents such as micronutrients, enzymes, vitamins and growth stimulators. SEAGROWING stimulates enzymatic processes acting as natural catalyst to promote balanced growth enabling the crop to fulfil its genetic potential. SEAGROWING is rapidly absorbed by foliar surfaces and moves into the cells increasing metabolic activity. SEAGROWING can be used both as a foliar application and in fertigation systems.

DOSAGE AND USAGE PROCEDURES

Citrus Top Fruit	150 g/Hl (foliar) 5 Kg/Ha (fertigation)
Grapes	100 g/Hl (foliar) 5 Kg/Ha (fertigation)
Field and Greenhouse vegetables	100 g/Hl (foliar) 5 Kg/Ha (fertigation)
Floriculture	100 g/Hl (foliar) 5 Kg/Ha (fertigation)

According to the DPR 26.10.1972, n 633, for SEAGROWING has to be applied a 10% VAT.



Dosage always depends on different factors (plant's age, growth stage, climatic conditions, temperature, etc) and can be increased or decreased according to your requirements. 5-6 application are advised during the whole vegetative cycle. IN CASE OF FOLIAR APPLICATION, MIXTURES WITH COPPER BASED PRODUCTS TO BE AVOIDED.

STIMOLEAF **BIOSTIMULANT**

ORGANIC FLUID FERTILIZER

STIMOLEAF is a natural bio-stimulant high in amino-acid content derived from the hydrolysis of animal proteins. STIMOLEAF is formulated to increase vegetative growth, especially applicable after unfavorable climatic conditions (frosts, strong winds, hail, etc.)

DOSAGE AND USAGE PROCEDURES

Citrus - Fruit trees	300 g/Hl (foliar)
Grapes	10 Kg/Ha (fertigation)
Open Field vegetables	250 g/Hl (foliar) 10 Kg/Ha (fertigation)
Greenhouse vegetables	300 g/Hl (foliar)
Floriculture	10 Kg/Ha (fertigation)



COMPOSITION

Organic Nitrogen (N)8,20% water soluble
Organic Carbon (C)24,65%
C/N Ratio2,7
Average molecular weight1500 dalton
Glycine Ratio1,1 (Proline + Hydroxyproline)
Hydrolysis on dry matter>330
Free Aminoacids>10%





Bottles Kg. 1

TO BE AVOIDED.

Tanks Ka. 6

